



**Aotearoa New Zealand national
paediatric early warning system
and paediatric vital signs chart
User guide**

March 2023

Contents

Document purpose.....	3
1. Introduction and scope	4
2. Chart overview.....	4
2.1. Recording vital signs: scoring area.....	6
2.2. Recording vital signs: non-scoring area	7
2.3. Mandatory escalation pathway	7
2.4. Modifications box.....	7
2.5. National and local tools	7
3. Clinical use of the PVSC	9
3.1. Documenting scoring vital signs	9
3.2. Documenting vital signs that do not contribute to the PEWS.....	16
3.3. Partial PEWS.....	20
3.4. Accountability	20
3.5. Modifying PEWS triggers	20
3.6. Calculating the total score and using single-parameter triggers.....	22
3.7. Escalating care.....	27
4. Design and printing information	28
4.1. Required amendments.....	28
4.2. Allowable amendments	28
4.3. Print specifications	29
4.4. Colour specifications	30

Figures

Figure 1: Overview of the front of the paediatric vital signs chart.....	5
Figure 2: Overview of the back of the paediatric vital signs chart	5
Figure 3: Paediatric early warning score colour key	6
Figure 4: Respiratory rate (example from 12+ years PVSC)	9
Figure 5: Assessment of respiratory distress guide from the back of the PVSC.....	10
Figure 6: Example of documenting respiratory distress.....	10
Figure 7: Oxygen	11
Figure 8: Respiratory support mode key as it appears on the back of all four PVSC	11
Figure 9: Examples of oxygen documentation	12
Figure 10: CPAP example	13
Figure 11: Oxygen saturation examples.....	14
Figure 12: Heart rate examples.....	14
Figure 13: Example of documentation of a central capillary refill of 3 or more seconds.....	15

Figure 14: Blood pressure examples.....	15
Figure 15: Level of consciousness examples	16
Figure 16: Temperature example	17
Figure 17: Assessment of pain tools on each chart	18
Figure 18: PEW score recorded from an incomplete set of observations	20
Figure 19: Initialling the chart	20
Figure 20: Example of modification due to chronic condition	21
Figure 21: Example of modification for a teenager with athletic bradycardia	21
Figure 22: Space to indicate if tamariki have an end-of-life pathway plan	21
Figure 23: Escalation pathway	27
Figure 24: Z-fold demonstrated with the PVSC.....	29

Tables

Table 1: PEWS score matrix	23
Table 2: Allowable amendments to the PVSC	28
Table 3: Colour specifications for the PVSC	30

Published March 2023 by the Health Quality & Safety Commission,
PO Box 25496, Wellington, New Zealand.



Published under Creative Commons License CC BY-NC 4.0. This means you may share and adapt the material provided you give the appropriate credit, provide a link to the licence, and indicate if changes were made. You may not use the material for commercial use. For more information about this licence, see: <https://creativecommons.org/licenses/by-nc/4.0/>

Document available online at: www.hqsc.govt.nz

Enquiries to: info@hqsc.govt.nz

**Te Kāwanatanga
o Aotearoa**
New Zealand Government

Document purpose

This user guide is for any health professional implementing and improving paediatric early warning systems within Aotearoa New Zealand hospitals. It explains how to complete and use the draft paediatric vital signs chart.

1. Introduction and scope

The national paediatric vital signs chart (PVSC) and paediatric early warning system (PEWS) are tools for detecting clinical deterioration in tamariki receiving hospital care in Aotearoa New Zealand. This user guide explains how to complete and use the PVSC correctly.

Begin the PVSC for any tamariki who is assessed as requiring measurement and recording of vital signs.

Tamariki physiology alters as they age, impacting on the normal ranges for vital signs. As such, there are four age-based PVSCs, which must be used for the correct age group:

- 0–11 months
- 1–4 years
- 5–11 years
- 12+ years.

Tamariki who need to receive care in an intensive care unit or high-dependency unit may not require a PVSC because these units usually have their own specific charts. The PVSC can be used in these settings; however, for consistency, the same parameters and scoring cut-offs for each vital sign should be used.

Before tamariki are transferred from an intensive care unit or high-dependency unit to a ward area, chart the most recent set of vital signs on an age-appropriate PVSC and include a plan to address any ongoing abnormalities in a set timeframe.

The total paediatric early warning score (PEW score) is calculated using measurements taken across seven vital sign parameters. The score increases as the vital signs deviate further from the normal zone. The total score triggers an escalating clinical response, so clinicians with the right skills can intervene and manage any deterioration promptly.

The Health Quality & Safety Commission developed the PVSC and PEWS based on the best available human factors and clinical evidence.

2. Chart overview

The PVSC is a two-sided document, printed front and back. Figure 1 and Figure 2 illustrate the main areas of both sides. The four age-based PVSCs all have the same layout. The vital sign parameter ranges differ by age, as does some of the content on the back.

This section provides a brief overview of each chart area. For more detail about the clinical use of the PVSC, see [Section 3](#) of this guide.

Figure 1: Overview of the front of the paediatric vital signs chart

The front of the paediatric vital signs chart is divided into several key sections:

- Recording vital signs: scoring area:** The top half of the chart, containing tables for Respiratory Rate, Respiratory Distress, Oxygen, Heart Rate, Central Capillary Refill, Blood Pressure, and PEWS TOTAL. A red arrow points from the PEWS TOTAL row to the modifications box.
- Recording vital signs: non-scoring area:** The bottom half of the chart, containing tables for Whānau concern, Consciousness, Temperature, and Pain Score.
- Mandatory escalation pathway:** A vertical column on the right side of the chart, showing PEWS 1-3, PEWS 4-5, PEWS 6-7, and PEWS 8+.
- Modifications box:** A table on the right side of the chart, used for recording modifications to PEWS scores.

Figure 2: Overview of the back of the paediatric vital signs chart

The back of the paediatric vital signs chart includes the following sections:

- Local tools:** A section on the left side of the chart, containing tables for Whānau concern, Consciousness, Temperature, and Pain Score.
- National tools:** A section on the right side of the chart, containing tables for Revised FLACC observational pain tool, Assessment of respiratory distress guide, and Respiratory support mode.
- Faces pain scale - revised:** A scale at the bottom of the chart, showing six faces with increasing expressions of pain, from 0 to 10.

2.1. Recording vital signs: scoring area

The purpose of the scoring area is to document the vital sign measurements of the core parameters that contribute to the total PEW score. The vital signs included for scoring are based on Bedside PEWS.¹ These seven parameters are:

- respiratory rate
- respiratory distress
- oxygen
- oxygen saturation
- heart rate
- central capillary refill
- blood pressure.

Where the observation plots on the chart determines its score: 0,1, 2 or 4. The score increases as the observation becomes more abnormal.

When documentation in the scoring area is done in a clear and consistent manner, it is easy to detect trends indicating improvement or deterioration. It is easier to visualise trends if you record using a symbol such as an 'X' rather than writing a numerical value (the exception being the oxygen saturation parameters, where the numerical value is marked on the chart). However, if a measured value falls outside the range able to be plotted on the chart, it can still be recorded by writing the numerical value at the upper or lower margin of the chart area for that parameter.

Scores for the individual parameters are then added to calculate the total PEW score, which is documented in the bottom row of the scoring area. Abnormal vital signs associated with a given score are also identified by the differently coloured zones on the chart. Individual vital sign parameters in blue zones trigger escalation actions that the escalation pathway sets out, **regardless of the total score.**

The total PEW score reflects the sum of the scores from seven vital signs. As it is not always possible or necessary to obtain a complete set of observations, for example, blood pressure in distressed tamariki, there needs to be a way of indicating that an incomplete set of observations has been used to generate a score. The convention on this chart is to add a plus (+) to the value recorded in the PEWS Total box, eg, 3+. See [Section 3.3](#) for more detail.

Figure 3: Paediatric early warning score colour key

Paediatric early warning score colour key				
0	1	2	4	RRT: rapid response team

¹ Parshuram CS, Hutchison J, Middaugh K. 2009. Development and initial validation of the Bedside Paediatric Early Warning System score. *Crit Care* 13: R135. URL: <https://doi.org/10.1186/cc7998>.

2.2. Recording vital signs: non-scoring area

The purpose of the non-scoring area is to document the vital sign parameters that do not contribute to the total PEW score. These parameters are:

- whānau concern
- level of consciousness
- temperature
- pain score.

There is also space for the staff member recording the observations to enter their initials. More detailed documentation in the clinical notes will include name and designation.

2.3. Mandatory escalation pathway

The escalation pathway states the actions to be taken when a particular total PEW score is reached or a blue zone observation occurs. An increasing total PEW score indicates a higher risk of deterioration. Each organisation defines for their own location what the response to each level of the escalation pathway will be, reflecting available resources and processes of care. This includes determining the correct emergency response terminology for the blue zone, for example, rapid response team (RRT).

2.4. Modifications box

The modifications box is used to individualise escalation triggers for tamariki with chronic disease or vital sign abnormalities that do not necessarily represent clinical deterioration. Modifications are usually only required for one vital sign parameter but can be made to several. They can be made for the duration of admission or can be time limited. They should be made after considering the potential risks associated with altering the score an abnormal parameter will generate. The process for making modifications, including who can make them, is defined locally.

For example, tamariki with a known cardiac condition resulting in lower oxygen saturation levels may inappropriately trigger escalation of care unless a single parameter modification for oxygen saturation is made.

Caution: This area is not to be used to modify an overall score; it is only for individual vital sign parameters. See Section 3.5 for more information.

2.5. National and local tools

The back of the PVSC is divided into two sections: national and local tools.

For sites using an electronic patient observation system, the method of accessing these tools needs to be locally determined if they cannot be incorporated into the electronic system. For example, this may include posters in the clinical environment or laminated cards at computer stations.

The national tools are a non-editable set of resources to support the consistent use, completion and interpretation of vital signs on the front of the chart. The national tools are:

- assessment of respiratory distress guide
- respiratory support mode abbreviations
- pain scores.

The pain scoring tools vary between age-banded charts to reflect the appropriate tools to use for different age groups.

The local tools section is editable, and the content is defined locally. It can be used to document non-scoring observations such as weight, blood sugar level, phlebitis score or pressure injury assessment. Associated guidance that explains how these observations are defined may be included. The multidisciplinary team will need to consider carefully what to include, if anything, in this space; it must be clinically relevant and enhance the care provided to tamariki. If you are adding elements to this section, you will need support from your local communications team or design team.

3. Clinical use of the PVSC

This section describes clinical use of the PVSC and includes:

- documenting vital signs
- modifying PEWS triggers
- calculating PEW scores
- mandatory escalation of care.

3.1. Documenting scoring vital signs

Instructions for documenting each vital sign parameter, the accepted abbreviation and the unit of measurement appear in the grey column to the left of the scoring area of the PVSC. Where measurements exceed the range on the chart, write the numerical value on the chart at the upper or lower margin of the chart area for that parameter.

Date and time

The top two rows of the PVSC provide space for recording the date and time that each set of observations was taken. Use the 24-hour clock format for the time. Be aware that the columns are evenly spaced across the chart but the time interval between observations may vary.

Respiratory rate

Each row of the scoring area for respiratory rate (Figure 4) represents a range associated with varying severity of abnormality. Note that the vital sign parameters vary between charts, so pay careful attention to ensure the correct score is generated.

Figure 4: Respiratory rate (example from 12+ years PVSC)

Respiratory Rate (breaths/min) <i>mark RR with X</i>	≥ 35		Respiratory Rate (breaths/min) <i>mark RR with X</i>	≥ 35	
	30-34			30-34	
	25-29			25-29	X
	20-24			20-24	
	15-19	X		15-19	
	12-14			12-14	
	10-11			10-11	
	5-9			5-9	
	≤ 4			≤ 4	

Respiratory rate = 18 breaths/minute, white zone, score 0 Respiratory rate = 27 breaths/minute, yellow zone, score 1

Respiratory distress

Assess the level of respiratory distress using the 'Assessment of respiratory distress guide' on the back of the chart (Figure 5), which is based on a tool developed by the Royal

Children’s Hospital Melbourne.² It is the same across all four age-based PVSC and is scored at the level of the most severe sign.

Figure 5: Assessment of respiratory distress guide from the back of the PVSC

Assessment of respiratory distress guide			
	Mild	Moderate	Severe
Airway	<ul style="list-style-type: none"> • Stridor on exertion or crying • Wheeze present 	<ul style="list-style-type: none"> • Some stridor at rest • Wheeze marked 	<ul style="list-style-type: none"> • Stridor at rest • New onset of stridor • Wheeze severe • Silent chest
Behaviour and feeding	<ul style="list-style-type: none"> • Normal • Talks in sentences 	<ul style="list-style-type: none"> • Some or intermittent irritability • Difficulty talking or crying • Difficulty feeding or eating 	<ul style="list-style-type: none"> • Increased irritability and/or lethargy • Looks exhausted • Unable to talk or cry • Unable to feed or eat
Accessory muscle use	<ul style="list-style-type: none"> • Mild intercostal and suprasternal recession 	<ul style="list-style-type: none"> • Moderate intercostal and suprasternal recession • Tracheal tug • Nasal flaring • Head bobbing 	<ul style="list-style-type: none"> • Marked intercostal and suprasternal recession
Other		<ul style="list-style-type: none"> • May have brief apnoea 	<ul style="list-style-type: none"> • Gaspings, grunting • Extreme pallor, cyanosis • Increasingly frequent or prolonged apnoea
Score at the level of severest sign. Note that not all features are relevant to all conditions.			

The level of respiratory distress can then be marked with an ‘X’ in the appropriate box (Figure 6). There are four levels of respiratory distress: nil, mild, moderate and severe.

Figure 6: Example of documenting respiratory distress

Respiratory Distress <i>mark RD with X</i>	Severe	
	Moderate	X
	Mild	
	Nil	

Oxygen

Tamariki receive a variety of oxygen therapies and modes of respiratory support. The challenge is to provide a format that captures information correctly and accurately reflects

² The Royal Children’s Hospital Melbourne. 2019. *Observation and continuous monitoring* (Clinical Guidelines (Nursing)). Melbourne: The Royal Children’s Hospital. URL: www.rch.org.au/rchcpg/hospital_clinical_guideline_index/Observation_and_continuous_monitoring/ (accessed 27 July 2021).

the oxygen therapy in use. The oxygen score is based mainly on the FiO₂, where an increasing oxygen requirement indicates a deterioration.

In some tamariki, for example those receiving nasal prong oxygen, the flow rate is varied but the FiO₂ remains unchanged at 100%. In another example, tamariki may have high-flow oxygen therapy in place, in which case the flow rate is stable and the FiO₂ is varied. This information can be recorded on the PVSC to generate a score for the oxygen parameter, and by recording the numerical value on the chart, the trend in the amount of respiratory support needed is visible. Tamariki on room air only are marked with an 'X' (Figure 7). Room air refers to no respiratory support at all.

Figure 7: Oxygen

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	
	< 4L or < 35%	
	Room air X	X
	Mode	
	High flow rate	

Record the mode of oxygen delivery using the abbreviations from the 'Respiratory support mode' panel on the back of the chart (Figure 8).

Figure 8: Respiratory support mode key as it appears on the back of all four PVSC

Respiratory support mode		
NP = nasal prongs	M = face mask	HF = high flow
R = non-rebreather mask	C = CPAP	B = BiPAP
TH = tracheostomy humidification	HO ₂ = humidified oxygen	

When tamariki are receiving high-flow oxygen therapy, the weight-based flow rate is recorded in the high-flow rate row. This allows the correct flow rate to be confirmed. The numerical value of the FiO₂ being delivered while on high-flow therapy can be recorded in either one of the two top boxes in the oxygen section, as appropriate.

We have not allocated space to record the pressure (in cmH₂O) being delivered to tamariki on CPAP (continuous positive airway pressure), although know that this may be varied during treatment. This is because it is uncommon for tamariki to receive CPAP outside a higher acuity area, where there is likely to be a different chart in use, and also due to space constraints on the chart.

Examples of different oxygen therapies and how to record them are shown in Figure 9.

Figure 9: Examples of oxygen documentation

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	
	< 4L or < 35%	
	Room air X	x
	Mode	
	High flow rate	

Tamariki on no respiratory support.

This parameter scores 0.

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	
	< 4L or < 35%	1L
	Room air X	
	Mode	NP
	High flow rate	

Tamariki on 1 L/min of oxygen via nasal prongs.

This parameter scores 2.

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	
	< 4L or < 35%	28%
	Room air X	
	Mode	HF
	High flow rate	14L

Tamariki weighing 7 kg on high-flow therapy with an FiO₂ requirement of 28%.

This parameter scores 2.

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	
	< 4L or < 35%	21%
	Room air X	
	Mode	HF
	High flow rate	14L

Tamariki weighing 7 kg on high-flow therapy with an FiO₂ requirement of 21%.

This parameter scores 2.

Please note that 21% in this context is not referring to room air. The tamariki is on high flow therapy and requires respiratory support.

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	5L
	< 4L or < 35%	
	Room air	X
	Mode	M
	High flow rate	

Tamariki receiving 5 L/min of oxygen via a mask.

This parameter scores 4.

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	
	< 4L or < 35%	0.5L
	Room air	X
	Mode	H ₂ O ₂
	High flow rate	

Tamariki receiving 0.5 L/min of humidified oxygen.

This parameter scores 2

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	36%
	< 4L or < 35%	
	Room air	X
	Mode	HF
	High flow rate	20L

Tamariki weighing 10 kg on high-flow therapy with an FiO₂ requirement of 36%.

This parameter scores 4.

Continuous positive airway pressure (CPAP)

If tamariki are receiving CPAP in the ward setting, the PVSC can be adapted to record settings and generate a score. Do this by changing the name of some of the rows to reflect what you need to record on CPAP.

CPAP generates a score based on the FiO₂. Record this on the chart in the same way you would for tamariki on high flow.

The positive end expiratory pressure (PEEP) is measured in cmH₂O and can be recorded in the 'mode' row. Cross out 'mode' and write 'Pressure cmH₂O'.

The CPAP flow is measured in litres per minute and can be recorded in the 'high flow rate' row. Cross out 'high' and write CPAP (see Figure 10).

Figure 10: CPAP example

Oxygen (L/min or FiO ₂ %) <i>write value</i>	≥ 4L or ≥ 35%	35%	35%
	< 4L or < 35%		
	Room air	X	
	Pressure cm H ₂ O Mode	5	8
	CPAP High flow rate	10	10

Oxygen saturation

Document the numerical value for oxygen saturation in the relevant box of the scoring area (Figure 11).

Figure 11: Oxygen saturation examples

Oxygen Saturation (%) <i>write SpO₂</i>	≥ 95	96	Oxygen saturation = 96% Scores 0
	91-94		
	≤ 90		

Oxygen Saturation (%) <i>write SpO₂</i>	≥ 95		Oxygen saturation = 93% Scores 1
	91-94	93	
	≤ 90		

Heart rate

Document heart rate by placing an 'X' in the relevant box of the scoring area (Figure 12). Each row of the scoring area for heart rate corresponds to a numerical range of 10 (eg, a heart rate in the 70s, 80s, 90s). In this way, you can clearly identify the relevant coloured zone if the heart rate value falls exactly on the line between zones (eg, a heart rate of 80 beats per minute is within the 80s range so scores within the yellow zone).

Figure 12: Heart rate examples

Heart Rate (bpm) <i>mark HR with X</i> <i>write value if off scale</i>	≥ 180		Heart Rate (bpm) <i>mark HR with X</i> <i>write value if off scale</i>
	170s		
	160s		
	150s		
	140s		
	130s		
	120s		
	110s		
	100s		
	90s	X	
	80s		
	70s		
	60s		
≤ 59			

Heart rate = 96, white zone, score 0

Heart rate = 80bpm, yellow zone, score 1

Central capillary refill

When looking at skin perfusion as a marker of cardiovascular status, the most accurate assessment is made by determining central capillary refill.³ This is obtained by pressing a finger on the central chest, holding for 5 seconds and releasing. The white blanched area should fully regain colour in under 3 seconds. The observation is marked with an 'X' according to the time taken (Figure 13).

³ Royal College of Nursing. 2017. Standards for assessing, measuring and monitoring vital signs in infants, children and young people. London: Royal College of Nursing.

Figure 13: Example of documentation of a central capillary refill of 3 or more seconds

Central Capillary Refill <i>mark CR with X</i>	≥ 3 sec	X
	< 3 sec	


Blood pressure

The systolic blood pressure is the measurement used to generate a score for this vital sign. Document both systolic and diastolic pressure on the chart as this is important clinical information.

Measure blood pressure using an appropriate size of cuff and document the cuff size on their care plan.

Each row of the scoring area for blood pressure corresponds to a numerical range of 10 (eg, a systolic blood pressure in the 90s, 100s, 110s). At times, coloured zones change in the middle of a row as in Figure 14, where the zone change is 125. That is, 124 is in the yellow and 125 is in the orange. If the observation plots on the line, score according to the colour above.

Figure 14: Blood pressure examples



BP = 105/70, white zone, score 0

BP = 125/75, orange zone, score 2

3.2. Documenting vital signs that do not contribute to the PEWS

Whānau concern

Y = Yes N = No A = away/asleep/unavailable

This section requires staff to check in with whānau as to whether they have any concerns about the condition of their tamariki. Asking regularly gives whānau the opportunity to speak up and voice concerns. Sharing the PVSC with whānau as part of the admission and orientation process, and when completing observations, supports communication, helps understanding and encourages partnership.

If there is concern, mark with a **Y** on the chart. Escalation of this concern can occur, even if the other observations and the total PEW score are normal. Staff may find that, with listening to the concern, explanation or further discussion alleviates it. Document detail in the progress notes, including any action taken because of the concern being raised.

If there is no concern, mark with an **N**.

If whānau members are away, asleep or unavailable, mark with an **A**. This can then act as a reminder for other staff to check in with whānau when they are next available.

The whānau concern section of the PVSC needs to link in with local organisation policy regarding Kōrero mai⁴ processes to support patient, family and whānau escalation of concerns.

Level of consciousness

Assess level of consciousness using the alert, voice, pain or unresponsive (AVPU) scale. Document your assessment by placing an 'X' in the relevant box of the scoring area (Figure 15). If tamariki are asleep, this can also be noted. Once you have assessed tamariki and determined them to be sleeping normally, annotate 'Asleep' on the chart within the alert section of 'Level of consciousness'. An example of this is when you enter the room you see a baby stir to the noise but not wake fully. It is important to assess tamariki closely and not assume they are sleeping.

Figure 15: Level of consciousness examples

Level Of Consciousness <i>mark LOC with X</i>	Alert	X	Level Of Consciousness <i>mark LOC with X</i>	Alert	Asleep
	Voice			Voice	
	Pain			Pain	
	Unresponsive			Unresponsive	

Temperature

Measure temperature using a consistent method each time (eg, oral, axillary, tympanic), and record the method in the care plan. Document temperature by placing an 'X' in the relevant position on the chart. Figure 16 shows the documentation of 38.5°C. You may also choose to write the actual value, to one decimal place, on the chart, but if doing so always write it

⁴ Health Quality & Safety Commission. Kōrero mai – patient, family and whānau escalation. Wellington: Health Quality & Safety Commission. URL: www.hqsc.govt.nz/our-programmes/patient-deterioration/workstreams/patient-family-and-whanau-escalation/.

above the X. For temperatures that are outside the plottable area, write the actual value, to one decimal place, on the chart.

Figure 16: Temperature example

Temperature (°C) <i>mark Temp with X</i> <i>write value if off scale</i>	≥ 40			
	39s			
	38s	X		
	37s			
	36s			
	≤ 35			

Pain score

Record pain on the PVSC to help interpret abnormal vital signs and manage pain effectively.

The PVSC includes three pain scoring tools: the Numerical rating scale, Faces pain scale – revised⁵ and the revised FLACC observation pain tool.⁶ The score is documented at rest and with movement.

Many clinicians will be familiar with the FLACC observational pain tool, but note the **revised** version is included here. This is an objective tool that has been validated for use in non-verbal and cognitively impaired tamariki. Therefore, it can be used both for a 4-month-old baby and 7-year-old tamariki with global developmental delay, for example.

The tool used depends in part on age of the tamariki (Figure 17). The age reflects the required cognitive ability of tamariki to be able to use the numeric and Faces tool. The tool being used can be documented in their care plan.

⁵ Hicks CL, von Baeyer CL, Spafford P, et al. 2001. The Faces Pain Scale – Revised: Towards a common metric in pediatric pain measurement. *PAIN* 93: 173–83.

⁶ Malviya S, Vopel-Lewis T, Burke C, et al. 2006. The revised FLACC observational pain tool: improved reliability and validity for pain assessment in children with cognitive impairment. *Pediatric Anaesthesia* 16: 258–65.

Figure 17: Assessment of pain tools on each chart


Age-based PVSC	Numerical	Faces pain scale	Revised FLACC
0–11 months	No	No	Yes
1–4 years	No	Yes	Yes
5–11 years	Yes	Yes	Yes
12+ years	Yes	Yes	Yes


Numerical rating scale


No pain |-----| Worst pain
 0 1 2 3 4 5 6 7 8 9 10


On a scale of 0 -10, with 0 being no pain and 10 being the worst pain you can imagine, what number are you feeling right now on movement and at rest?


Faces pain scale - revised



0


2


4


6


8


10

Can you point to the face that shows how much you hurt?

Revised FLACC observational pain tool			
Categories	Scoring		
	0	1	2
Face	No expression or smile	Occasional grimace or frown, withdrawn, disinterested; appears sad or worried	Frequent to constant frown, clenched jaw, quivering chin; <i>distressed looking face; expression of fright or panic</i>
			Individualised behaviour described by family:
Legs	Normal position or relaxed; usual muscle tone and motion to arms and legs	Uneasy, restless, tense; occasional tremors	Kicking, or legs drawn up; <i>marked increase in spasticity; constant tremors or jerking</i>
			Individualised behaviour described by family:
Activity	Lying quietly, normal position, moves easily; regular rhythmic breaths (respiration)	Squirming, shifting back and forth, tense or guarded movements; mildly agitated (head back and forth, aggression); shallow, splinting breaths (respirations); occasional sighs	Arches, rigid, or jerking; <i>severe agitation; head banging; shivering (not rigors); breath holding, gasping, or sharp intake of breaths; severe splinting</i>
			Individualised behaviour described by family:
Cry	No cry (awake or asleep)	Moans or whimpers, occasional complaint; occasional verbal outburst or grunt	Crying steadily, screams or sobs, frequent complaints; <i>repeated outbursts; constant grunting</i>
			Individualised behaviour described by family:
Consolability	Content, relaxed	Reassured by occasional touching, hugging, or 'talking to'; can be distracted	Difficult to console or comfort; <i>pushing away caregiver; resisting care or comfort measures</i>
			Individualised behaviour described by family:
Rate the child in each of the five measurement categories, add together, and document total pain score (0 – 10).			
Children who are awake:	Observe for at least 1-2 minutes. Observe legs and body uncovered. Reposition child or observe activity, assess body for tenseness and tone. Initiate consoling interventions if needed.		
Children who are asleep:	Observe for at least 2 minutes or longer. Observe legs and body uncovered. If possible, reposition the child. Touch the body and assess for tenseness and tone.		
This tool can be used for all non-verbal children. The additional descriptors (in italics) are validated in children with cognitive impairment. The nurse can review with parents/caregivers the descriptors within each category. Ask the parents/caregivers if there are additional behaviours that are better indicators of their child experiencing pain. Add these behaviours to the tool in the appropriate category.			

3.3 Partial PEWS

Some sets of observations will not include measurement of all the parameters required to calculate a total PEW score. The risk of not indicating that a score is based on an incomplete set of observations is that the user may underestimate how unwell tamariki are and use an escalation pathway lower than what is required.

In this situation, the PEW score from an incomplete set of observations is indicated by a plus (+) (Figure 18), and the user should act on the escalation pathway mandated by this score. They should also consider whether any other actions need to be taken, including completing the full set of observations, and follow local monitoring policy on what observations are required for specific circumstances.

Multiple sets of partial observations may occur in certain circumstances, for example, taking a series of blood pressure measurements after adjustment of medication. However, this practice should be the exception rather than the rule, to maintain the value of the PEW score

Figure 18: PEW score recorded from an incomplete set of observations



3.4. Accountability

To fulfil documentation requirements, the clinician who undertook the observations must place their initials at the bottom of the chart (Figure 19).

Figure 19: Initialling the chart

Pain Score <i>write score (0-10)</i>	Rest	1	3
	Movement		
Initials		AL	

3.5. Modifying PEWS triggers

It is possible to modify the PEWS triggers for tamariki when chronic disease or abnormal vital sign parameters are present that do not necessarily represent clinical deterioration. This section of the chart is to capture what is 'normally abnormal'. Individual vital sign parameters are altered, not total PEW scores.

When making modifications, consider the clinical risk to tamariki of normalising an abnormal vital sign. This risk is mitigated by discussing modifications with a senior clinician and reviewing them at regular intervals, so they remain appropriate as the condition of tamariki changes. Document the duration for which modifications apply to ensure there is a timely clinical review.

Local policy on modifying PEWS triggers should be in place to ensure practice is appropriate for your local population and hospital context. This includes identifying which clinicians can make a modification and how often modifications need to be reviewed. Please ensure that clinicians tasked with making modifications have received appropriate education.

When a new PVSC is used, the modifications will need to be reviewed, redocumented and signed if they are still applicable. Staff should not just assume modifications remain in place. Figure 20 and Figure 21 give examples of appropriate documentation.

Note: If tamariki are generating high PEWS scores, the scores are likely to remain high until treatment takes effect. In this situation, do not use the modification section to stop escalation. Rather, the review generated by the high PEWS score should result in an understanding of why the PEWS score is elevated and clear documentation in the patient notes of a treatment plan. This should include specific signs of further deterioration that should be notified, who should be called and a recommended time for the next review.

Having a modification in place should not stop staff calling a clinical emergency if a parameter enters the blue zone or if staff feel a significant deterioration is imminent.

Figure 20: Example of modification due to chronic condition

Vital sign (use abbreviation)	Accepted values and modified PEWS	Date and time	Duration (hours)	Name and contact details
Oxygen saturation	$\geq 91\%$ score = 0 85–90% score = 1 $\leq 85\%$ score = 2	20/ 3 /21 11:30 am	Throughout admission	N. Rivera #6132
Reason: Normal saturations for child is 91% due to cyanotic congenital heart disease				

Figure 21: Example of modification for a teenager with athletic bradycardia

Vital sign (use abbreviation)	Accepted values and modified PEWS	Date and time	Duration (hours)	Name and contact details
HR	50–109 score = 0 40–49, score = 1 30–39, score = 2 < 30 score = 4	20/ 3 /21 11:30 am	until discharge	D. Ramoray #2611
Reason: Competitive rower, resting HR of 50 when well				

There is also a space to accommodate tamariki receiving palliative care. Underneath the modification section is the ability to identify if tamariki are on an end-of-life pathway (Figure 22). This section is editable to reflect the terminology and pathway that reflects the organisational policy.

Figure 22: Space to indicate if tamariki have an end-of-life pathway plan

END OF LIFE PATHWAY	<input type="checkbox"/>	/ /	:
---------------------	--------------------------	-----	---

3.6. Calculating the total score and using single-parameter triggers

Each scoring vital sign parameter has coloured zones (white, yellow, orange and red) that are associated with a score of 0, 1, 2 or 4, respectively. Add together the score for each of the seven vital sign parameters to get a total PEW score. The total score can trigger action according to the escalation pathway.

The PVSC has single parameter triggers, which are any observations falling in the blue zone. The blue zone is not associated with a score because any parameter in the blue zone indicates severe deterioration and requires an emergency response.

Table 1 illustrates the PEWS score matrix across the four charts.

Table 1: PEWS score matrix

PEWS matrix: 0–11 months

Score	RRT	4	2	1	0	1	2	4	RRT
Zone	Blue	Red	Orange	Yellow	White	Yellow	Orange	Red	Blue
Respiratory rate	≤ 9	10–19	20–24	25–29	30–49	50–54	55–69	≥ 70	
Respiratory distress					Nil	Mild	Moderate	Severe	
Oxygen					Room air		< 4 L/min or < HF 35%	≥ 4 L/min or ≥ HF 35%	
Oxygen saturation			≤ 90	91–94	≥ 95				
Heart rate	50–59	60–79	80–89	90–109	110–159	160–169	170–179	≥ 180	
Capillary refill time					< 3 sec			≥ 3 sec	
Blood pressure	≤ 49	50–54	55–64	65–74	75–99	100–119	120–149	≥ 150	

HF = high flow; RRT = rapid response team.

PEWS matrix: 1–4 years

Score	RRT	4	2	1	0	1	2	4	RRT
Zone	Blue	Red	Orange	Yellow	White	Yellow	Orange	Red	Blue
Respiratory rate	≤ 4	5–14	15–17	18–19	20–39	40–44	45–54	≥ 55	
Respiratory distress					Nil	Mild	Moderate	Severe	
Oxygen					Room air		< 4 L/min or < HF 35%	≥ 4 L/min or ≥ HF 35%	
Oxygen saturation			≤ 90	91–94	≥ 95				
Heart rate	50–59	60–69	70–79	80–89	90–139	140–149	150–159	≥ 160	
Capillary refill time					< 3 sec			≥ 3 sec	
Blood pressure	≤ 54	55–64	65–74	75–89	90–109	110–124	125–159	≥ 160	

HF = high flow; RRT = rapid response team.

PEWS matrix: 5–11 years

Score	RRT	4	2	1	0	1	2	4	RRT
Zone	Blue	Red	Orange	Yellow	White	Yellow	Orange	Red	Blue
Respiratory rate	≤ 4	5–11	12–14	15–19	20–29	30–34	35–44	≥ 45	
Respiratory distress					Nil	Mild	Moderate	Severe	
Oxygen					Room air		< 4 L/min or < HF 35%	≥ 4 L/min or ≥ HF 35%	
Oxygen saturation			≤ 90	91–94	≥ 95				
Heart rate	40–49	50–59	60–69	70–79	80–129	130–139	140–154	≥ 155	
Capillary refill time					< 3 sec			≥ 3 sec	
Blood pressure	≤ 54	55–69	70–79	80–89	90–119	120–139	140–169	≥ 170	

HF = high flow; RRT = rapid response team.

PEWS matrix: 12+ years

Score	RRT	4	2	1	0	1	2	4	RRT
Zone	Blue	Red	Orange	Yellow	White	Yellow	Orange	Red	Blue
Respiratory rate	≤ 4	5–9	10–11	12–14	15–24	25–29	30–34	≥ 35	
Respiratory distress					Nil	Mild	Moderate	Severe	
Oxygen					Room air		< 4 L/min or < HF 35%	≥ 4 L/min or ≥ HF 35%	
Oxygen saturation			≤ 90	91–94	≥ 95				
Heart rate	30–39	40–49	50–59	60–64	65–109	110–119	120–134	≥ 135	
Capillary refill time					< 3 sec			≥ 3 sec	
Blood pressure	≤ 64	65–69	70–84	85–99	100–134	135–149	150–189	≥ 190	

HF = high flow; RRT = rapid response team.

3.7. Escalating care

An increase in the total PEW score indicates the potential for or an actual deterioration in the condition of tamariki and warrants an escalation in their care. The PVSC displays locally determined mandatory escalation pathways depending on the total PEW score (Figure 23). However, care can be escalated regardless of the total PEW score, including activating an emergency response, if there is significant whānau or staff concern.

Figure 23: Escalation pathway

ESCALATE CARE FOR ANY PATIENT YOU OR THEIR WHĀNAU ARE WORRIED ABOUT, REGARDLESS OF VITAL SIGNS OR PEWS

Mandatory escalation pathway	
Total PEWS	Action
PEWS 1-3	
PEWS 4-5	
PEWS 6-7	
PEWS 8+	
Any vital sign in the blue zone	

4. Design and printing information

This section provides information about design and print requirements for the PVSC.

Your organisation needs to develop an escalation pathway appropriate to your local area before you can use it clinically. You can make a few other amendments according to locally agreed policy and practice. The details that follow cover:

- required amendments
- allowable amendments
- print specifications.

4.1. Required amendments

The escalation pathways should be developed with the clinicians using the PVSC and reflect local systems and practice. The pathway should use unambiguous language and briefly and clearly state expected actions and responses for each range of PEW score.

Enter the pathway in the 'Action' area of the chart (provided as an editable PDF). Use the guidance document, *Paediatric early warning system escalation mapping tool*,⁷ to develop the response for each level of physiological abnormality.

4.2. Allowable amendments

Table 2 sets out the amendments you may make to the PVSC.

Table 2: Allowable amendments to the PVSC

Chart area	Allowable amendment
Left-hand margin	For scanning purposes, you may add a barcode or QR code.
Left-hand margin	You may add a black-and-white version of the organisational logo below the patient label. Do not use coloured logos because they add visual clutter and distract from the main purpose of the chart.
Central column of the scoring area	You may replace 'RRT' with a locally relevant number or acronym, eg, '777'. Note, there is space for only three characters.
Beside 'Mandatory escalation pathway' heading	If you have different escalation pathways for different hospitals, you can add the name of the hospital next to this heading.
Right-hand bottom corner	You can replace 'End-of-life pathway' with your relevant local terminology.
Back of chart: 'Local tools'	This space can be edited to include locally defined parameters or tools. Do this in conjunction with your communications team or design team.

⁷ Available at: www.hqsc.govt.nz/our-programmes/patient-deterioration/publications-and-resources/publication/4350/.

4.3. Print specifications

The chart must be professionally printed to the following specifications:

- paper size A3 landscape
- double-sided (front and back)
- minimum paper quality 150 gsm (uncoated stock)
- fold: chart is creased twice at 205 mm and 310 mm (from the left-hand edge) and then
- Z-folded in front along these creases (Figure 24)
- hole-punched left-hand side with two holes
- no print offset required.

Figure 24: Z-fold demonstrated with the PVSC

Family Name: _____		Date of Birth: _____	Family Name: _____	
Given Name: _____		Gender: _____	Given Name: _____ Gender: _____	
NHf#: _____		AFFIX PATIENT LABEL HERE.		
Date of Birth: _____		Date of Birth: _____ NHf#: _____		
BE CARE FOR ANY PATIENT YOU OR THEIR WHĀNAU ARE D ABOUT, REGARDLESS OF VITAL SIGNS OR PEWS				
Respiratory escalation pathway				
EWS				
Action				
Respiratory Rate (breaths/min) mark RR with X	≥ 55			
	50-54			
Respiratory Distress mark RD with X	Severe	1-3		
	Moderate			
Oxygen (L/min or FiO ₂ %) write value	≥ 4L or ≥ 35%			
	< 4L or < 35%	4-5		
Oxygen Saturation (%) write SpO ₂	≥ 95			
	91-94			
Heart Rate (bpm)	≥ 180	6-7		
	170s			
Central Capillary Refill mark CR with X	≥ 3 sec			
	< 3 sec			
Blood Pressure (mmHg)	≥ 160			
	150s			
PEWS TOTAL	150s			
	140s			
Level Of Consciousness mark LOC with X	Unresponsive			
	≥ 41			
Temperature (°C)	40s			
	39s			
Pain Score write score (0-10)	38s			
	37s			
Initials	36s			
	Rest			
LIFE PATHWAY	Movement			

Modification to PEWS triggers

can be changed to prevent inappropriate escalation. All modifications made in line with hospital policy and regularly reviewed by the primary team. Query any modification that is not signed and dated.

Modification (signature)	Accepted values and modified PEWS	Date and time	Duration (hours)	Name and contact details
		/ /	:	

4.4. Colour specifications

CMYK (cyan, magenta, yellow and black) and RGB (red, green, blue) are two different ways of achieving a particular colour. Most printers print in CMYK, and most screens display RGB.

The PVSC was designed using CMYK colour because it is intended for print use. The CMYK colour values listed in Table 3 are used at 60 percent tint on the scoring area of the chart to enable clear documentation with a black or blue ballpoint pen. The colour values are used at 100 percent tint in the other sections of the chart, such as the columns to the right and left of the scoring area and the left-hand column of the escalation pathway.

If you want to use the colours in a screen display format, the CMYK values will need to be converted to RGB values (see Table 3). The CMYK and RGB colours have been used to shade the boxes in Table 3, and you will see there are minor differences. This is to be expected.

You can select RGB values in Word documents using the 'More colors' option in the shading or font colours tabs. Open 'More colors' then select 'Custom' and insert the RGB values.

Table 3: Colour specifications for the PVSC

Colours	C	M	Y	K	R	G	B
Yellow	0	0	40	0	255	255	153
Orange	0	12	27	0	255	224	186
Pink	0	25	0	0	255	192	216
Blue	18	9	0	0	209	232	255